



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : A61K 31/70, C07H 21/02, 21/04, C12N S10, 15/85		A1	(43) International Publication Date: 4 February 1999 (04.02.99)
(11) International Publication Number: WO 99/04800			

(21) International Application Number: PCT/US98/15130		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 22 July 1998 (22.07.98)		Published With international search report.	
(30) Priority Data: 08/898,094 22 July 1997 (22.07.97) US			
(71) Applicant (for all designated States except US): GENTRIX, LLC [US/US]; 160 Second Street, Cambridge, MA 02142 (US).			
(72) Inventors, and Inventors/Applicants (for US only): SEGAL, Andrew, H. [US/US]; 160 Second Street, Cambridge, MA 02142 (US). WILSON, Jeffrey [US/US]; 39 Burton Street, Brighton, MA 02135 (US).			
(74) Agent: WILLIAMS, Kathleen, M.; Banner & Wilcoff, Ltd., 28th floor, 28 State Street, Boston, MA 02109 (US).			

(54) Title: NUCLEIC ACID COMPOSITIONS AND METHODS OF INTRODUCING NUCLEIC ACIDS INTO CELLS

(57) Abstract

Bifunctional nucleic acid molecules are disclosed which can bind to a cell surface and which comprise a first nucleic acid which comprises an aptamer bonded to a second nucleic acid, the biological effector sequence, that possesses a biological activity. Also contemplated are templates, vectors and host cells comprising the bifunctional nucleic acid molecules and methods for introducing the biological effector sequence into an organism by administering a host cell transfected with the biological effector sequence.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China			PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
ER	Estonia	LR	Liberia	SG	Singapore		

# NUCLEIC ACID COMPOSITIONS AND METHODS OF INTRODUCING NUCLEIC ACIDS INTO CELLS

## Field of the Invention

The invention relates to the introduction of exogenous nucleic acid molecules into cells.

## Background of the Invention

Methods for the transfer of nucleic acids into cells are of great utility for both physicians and experimental biologists. For example, genes encoding proteins can be transferred to treat

diseases or to facilitate laboratory experiments. Furthermore, "antisense" nucleic acid molecules, which inhibit synthesis of a given polypeptide or polynucleotide by binding to the corresponding

template, can be similarly applied, as can polynucleotide enzymes ("ribozymes").

A number of methods have been developed to effect such transfer. In several such

methods, the nucleic acid molecule is physically complexed with a ligand that can bind to a

molecule on the surface of the target cells. The entire complex, including the nucleic acid, can

then be internalized by the cells via receptor-mediated endocytosis. To date, ligands used have

included monoclonal antibodies (see, e.g., Fertkol et al (1993) J Clin Invest 92:2394-400; Buschle

et al (1995) Hum Gene Ther 6:753-61), microboc-derived polypeptides (see, e.g., Ding et al

(1995) J Biol Chem 270:3667-76), and native ligands for targeted receptors (see, e.g., Sosnowski

et al (1996) J Biol Chem 271:33647-53; Christen and Roth (1996) Cancer Gene Ther 3:4-10).

Although nucleic acids have traditionally been conceived of as linear molecules that

encode information, it has recently been appreciated that some nucleic acid sequences adopt

conformations that allow them to bind to other molecules, such as polypeptides, specifically and

with high affinity via non-Watson-Crick interactions. Nucleic acid molecules with this property

are known as "nucleic acid ligands".

Nucleic acid ligands consisting of RNA, single-stranded DNA, and double-stranded DNA

have been described. Little is known about the structural features that determine the binding

capacity and specificity of such ligands. Nevertheless, it is clear that a proper secondary

structure is indispensable. "Secondary structure" refers to stable structural features formed by

30 intrastand Watson-Crick base pairing. Specific types of secondary structural motifs, defined by their internal juxtapositions of paired and unpaired segments, include, e.g., motifs conventionally known as hairpin loops, symmetric bulged hairpin loops, asymmetric bulged hairpin loops, bulges, and pseudoknots (Schimmel (1989) Cell 58:9-12). The importance of secondary structure to the binding activity of nucleic acid ligands was illustrated, for example, by the studies of Uhlenbeck et al (1983) J Biomolec Structural Dynamics 1:539 and Romaniuk et al (1987) Biochemistry 26:1563.

35 Nucleic acid ligands have been used to bind soluble molecules and inhibit their biological activities (see, for example, Jellinek et al (1994) Biochemistry 33:10450-56; Bock et al (1992) Nature 355:564-66; Ishizaki et al (1996) Nat Med 2:1386-89). Nucleic acid ligands that bind to cell surface molecules are also possible (Gold and Tuerk, U.S. Pat. 5, 270,163; Gold and Tuerk, U.S. Pat. 5, 475, 096), and for the purposes of the invention are termed "aptamers".

40 It is the object of the invention to provide compositions and methods for introducing nucleic acids, including non-aptamer nucleic acids, into cells using aptamers as ligands. Specifically the invention pertains to bifunctional nucleic acid molecules that comprise an aptamer and another useful nucleic acid sequence that is not an aptamer.

### Summary of the Invention

45 One aspect of the invention features nucleic acid molecules, herein referred to as "bifunctional nucleic acid molecules", which can bind to a cell surface with at least micromolar affinity and which comprise: a first nucleic acid which comprises an aptamer bonded to a second nucleic acid that possesses a biological activity (herein referred to as a "biological effector sequence") and which is not an aptamer.

50 The invention thus relates to a nucleic acid molecule comprising a first nucleic acid sequence comprising an aptamer covalently linked to a second nucleic acid sequence comprising a biological effector sequence.

55 The invention also relates to a nucleic acid molecule comprising a first nucleic acid sequence comprising an aptamer linked via Watson-Crick base pairing to a second nucleic acid sequence comprising a biological effector sequence.

A molecule of the invention may also further comprise a third nucleic acid sequence

- which is an aptamer that is covalently linked to the nucleic acid molecule.
- A molecule of the invention may also further comprise a third nucleic acid sequence which is an aptamer that is linked via Watson-Crick base pairing to the nucleic acid molecule.
- In preferred embodiments, the third nucleic acid sequence comprises an aptamer that is different from the first nucleic acid sequence comprising an aptamer.
- Preferably, a molecule of the invention comprises DNA or RNA.
- The phrase "nucleic acid molecule" as used herein is intended to include structures that comprise nucleotides covalently bound to each other to form polymers that can, for example, be linear, cyclic, and/or branched. The bonds between sequential nucleotides in the polymer are referred to herein as a "backbone" of the molecule and can be, for example, phosphodiester, phosphorothioate, phosphoramidate, thioformacetate, carbamate, methylphosphonate, or peptide bonds. The molecules can be single-stranded and/or multi-stranded, but in a preferred embodiment are single- or double-stranded. According to the invention, a nucleic acid molecule can include naturally occurring and non-naturally occurring nucleotides, and can include non-nucleotide structures, e.g. amino acids, carbohydrates, and metal ions. The nucleotides may be chemically substituted at the ribose and/or phosphate and/or base positions. For example, the nucleotides can incorporate 5-pyr, 2'-amino, 2'-fluoro, or 2'-O-methyl groups. A nucleic acid molecule can include structures that are linked to each other by means other than covalent bonds, e.g. by Watson-Crick base pairing between overlapping complementary sequences.
- An "aptamer" is a nucleic acid molecule that is capable, by virtue of secondary and/or tertiary structure, of binding to a cell surface molecule such as a carbohydrate or a protein in a selective fashion, preferably with an affinity as strong as in the micromolar range (1-100 uM) and more preferably with an affinity even stronger, e.g., in the nanomolar range (1-100 nM affinity and 1-100pM affinity). That is, the aptamer will selectively bind to the target molecule or cell with an affinity that is at least 10-fold greater affinity than the affinity with which the aptamer binds to a non-target molecule. In general, an aptamer will comprise about 10-400 nucleotides, more preferably 20-100, and most preferably 25-50. "Selective" binding refers to specific binding to a target cell surface molecule but not to most other molecules on the cell surface or on other cells that do not contain the targeted cell surface molecule.
- Preferably, the aptamer and the biological effector sequence are linked via a covalent

- bond that is a phosphodiester bond.
- Where the aptamer and the biological effector sequence share a continuous backbone, a nucleotide sequence of the bifunctional nucleic acid molecule can contribute to the biological function of both.
- 90 In other preferred embodiments, the aptamer may be 3' to the biological effector sequence or 5' to the biological effector sequence. The aptamer and the biological effector sequence can be linked in a 3'-3' manner, e.g. via an acetal group. A bifunctional nucleic acid molecule can comprise multiple aptamers and/or biological effector sequences. In a preferred embodiment, the aptamer and biological effector sequence are admixed with a polycation, such as poly-L-lysine. In yet another preferred embodiment, the aptamer and the biological effector sequence are linked via a biotin-streptavidin interaction.
- 95 It is preferred that the aptamer of a bifunctional nucleic acid molecule be able to bind to a cell-surface polypeptide present on a cell, preferably with nanomolar to picomolar (1pM - 100 nM), preferably 10pM-50 nM, most preferably 50pM-10 nM) affinity.
- 100 In one embodiment, the biological effector sequence of a bifunctional nucleic acid molecule comprises a sequence that encodes a biologically or therapeutically useful polypeptide or polynucleotide. In a preferred embodiment, the sequence is double-stranded DNA and comprises a promoter that is operably linked to the coding sequence. In another preferred embodiment, the biological effector sequence is an mRNA, that is an RNA that translates to a polypeptide, which preferably comprises a 5' cap and a 3' poly-A sequence.
- 105 In another embodiment, the biological effector sequence comprises a sequence that is antisense to a nucleic acid that is present in the target cell. In preferred embodiments the antisense sequence can either 1) inhibit transcription of the target, or 2) inhibit reverse transcription of the target, or 3) inhibit translation of the target, or 4) inhibit replication of the target, or 5) inhibit the target from assuming a functional conformation.
- 110 In other preferred embodiments, the antisense sequence is complementary to at least five contiguous nucleotides of the target, or is complementary to at least seven contiguous nucleotides of the target, or is complementary to at least ten contiguous nucleotides of the target, or is complementary to at least fifteen contiguous nucleotides of the target, or is complementary to at least twenty contiguous nucleotides of the target. Thus, the antisense sequence may form a
- 115

Watson-Crick base-paired hybrid with a target RNA or DNA and inhibit expression (translation or transcription or regulatory binding) to the target molecule.

In another preferred embodiment, the biological effector sequence comprises a nucleic acid enzyme, e.g. a ribozyme.

120

In one embodiment, a bifunctional nucleic acid molecule can bind more than one cell surface molecule at once via its aptamers. In a preferred embodiment, the aptamer can concatenate and comprises first and second unpaired nucleic acids, each of which is complementary to itself, however, neither of which is complementary to the other. This structure allows concatemers of the aptamer to form and incorporate into the bifunctional nucleic acid molecule. A concatemer is at least two aptamers, and up to 1000 or even more (10,000) aptamers, but is preferably in the range of two-500 aptamers, and most preferably in the range of two - 50 aptamers. It is believed that such concatemers can cross-link the target receptors, increasing endocytosis.

125

The invention also pertains to vectors which include the bifunctional nucleic acid molecules of the invention, cells which have been introduced such vectors, and cells into which have been introduced the molecules of the invention.

130

In another aspect, the invention pertains to nucleic acid molecules that can be used in the production of bifunctional nucleic acid molecules. In a preferred embodiment, these molecules are DNA or RNA molecules that can serve as templates for the assembly of bifunctional nucleic acid molecules, e.g. by possessing Watson-Crick complementarity to a bifunctional nucleic acid molecule and by directing the assembly of a complementary nucleic acid, e.g. by the action of a nucleotide polymerase.

135

In another aspect, the invention pertains to compositions that include the nucleic acid molecules of the invention and that include substances that can facilitate the use of these nucleic acid molecules in vivo and/or in vitro. In one embodiment, the composition includes a pharmaceutically acceptable carrier.

140

In another embodiment, the composition includes a substance that improves the uptake of exogenous polynucleotides by cells, e.g. by inducing endocytosis and/or phagocytosis and/or macropinocytosis and/or micropinocytosis and/or potocytosis. In yet another embodiment, such compositions can include substances that disrupt intracellular vesicles

145

- containing the nucleic acid molecules of the invention.
- It is preferred that a composition of the invention comprise a bifunctional nucleic acid molecule and a substance that facilitates the intracellular trafficking of the biological effector sequence to its site of action. Such substances can include, for example, those that facilitate release from an intracellular vesicle or those that facilitate translocation to the nucleus.
- In yet another aspect, the invention pertains to methods of introducing biological effector sequences into cells.
- In a preferred embodiment, the complexes of the invention are admixed in vitro with cells bearing the receptor for the aptamer. It is further preferred that the cells are subjected to transfection with the complexes utilizing, e.g., calcium phosphate precipitation or electroporation. In another preferred embodiment, the cells are treated with an endosomal disruption agent. In another embodiment, the complexes of the invention are administered to an organism, e.g. an animal, a mammal, e.g. a primate, e.g. a human.
- In another aspect, the invention features a method of introducing a biological effector sequence into an organism, comprising introducing a biological effector sequence into a cell by contacting the bifunctional nucleic acid with a host cell, and administering the cell to an organism.
- Other features and advantages of the invention are described hereinbelow and in the claims.
- 165
- Detailed Description of the Invention**
- The invention is based on the recognition that a bifunctional nucleic acid, that is, a first nucleic acid sequence which is an aptamer specific for a cell surface molecule such as a protein or carbohydrate that is bound to a second nucleic acid sequence that is a biological effector sequence and not an aptamer can facilitate entry of the second nucleic acid sequence into a target cell. The bifunctional nucleic acid molecule is composed of two nucleic acid sequences or fragments that are bound together via covalently bonding or via Watson-Crick base-pairing along a portion of the two sequences, the portion being long enough to provide sufficient stability for the two molecules to form a hybrid and to facilitate entry of the second sequence into the target cell.
- 170
- 175



The contents of citations referred to herein are incorporated by reference in their entirety.

#### Production of Aptamers

An aptamer specific for a given target molecule is isolated from a heterogeneous library of polynucleotides by affinity purification against the target molecules. For example, in the

"SELEX" procedure described in Gold and Turk, U.S. Pat. 5, 270, 163, the following steps are performed:

1) The library is prepared as follows. For example, a random collection of oligonucleotides can be generated by programming an automated synthesizer to select a solution of multiple bases for incorporation at one or more steps. Ideally, this library contains at least about  $10^9$  nucleic acids, and no more than about  $10^{12}$ . If an RNA library is desired, it may be convenient to first obtain a corresponding DNA library and then, after cloning into a suitable plasmid, obtain the RNA by in vitro transcription.

2) The library is contacted with the target molecule under physiological chemical

conditions. Contacting can occur either with ligands and target free in solution, or with the target bound to a support. Examples of supports include agarose or sepharose matrices. The target can be coupled to such matrices either covalently or noncovalently.

3) The candidate aptamers with the highest affinities for the target (e.g., 5-50%

of the polynucleotides) are then partitioned from those with lower affinities. In one partitioning method, the target is a polypeptide and the candidate/target mixture is passed through a

nitrocellulose membrane and the membrane washed with, e.g., 200mM potassium acetate, 50mM Tris-HCl pH 7.7. This often elutes free candidates ("losers") while leaving aptamer/target complexes bound. The "winners" are then released from the target by washing the membrane with, e.g., 200ul 7M urea, 20mM Na citrate (pH 5.0), 1mM EDTA with 500 ul phenol (equilibrated with 0.1M Na acetate pH 5.2).

In another partitioning method (Toole et al, U.S. Pat. 5, 582, 981), free candidates are

eluted from a support-bound target using a suitable buffer. The target is then uncoupled from the support and the aptamer/target complexes collected. The nucleic acid molecules are isolated by standard denaturation techniques, such as phenol extraction.

If either of the above partitioning methods are used for a target that is a cell surface

molecule, it is preferred that only the extracellular portion of the molecule be employed in the

assay.

In a partitioning method that is useful for cell surface targets, cells are obtained that do

not express the desired target. The library is admixed with the cells, and the unbound candidates are collected. These candidates are then admixed with cells that are identical to those above,

except that they do express the target. For example, they may be transfected with a gene

encoding a polypeptide target, or enzymatically modified to bear a carbohydrate target. The cells

are spun down and the unbound candidates in the supernatant discarded. The cells are then

treated with phenol to release the aptamers.

4) The selected polynucleotides are amplified, typically by PCR. If the candidates are

RNA, they are reverse transcribed, the DNA is amplified, and RNA is again generated by reverse

transcription as above. The product is a library enriched for capacity to bind the target.

5) Steps 2-5 are repeated. In each round, binding and/or partitioning conditions can be

made more stringent. At least ten rounds are usually required for optimal results.

Cells that can bear targets for aptamers include bacterial, fungal, plant, yeast and

mammalian cells, e.g., malignant tumor cells, fibroblasts, endothelial cells, epithelial cells

including respiratory epithelial cells, pluripotent stem cells, leukocytes, endocrine cells such as

islet cells, hepatocytes, keratinocytes, melanocytes, pericytes, germ cells, neurons, myocytes

including cardiac myocytes, osteocytes, osteoblasts and chondrocytes.

Cell surface molecules that are candidate targets for aptamers include, e.g., the transferrin

receptor, the asialoglycoprotein receptor, the TSH receptor, FGF receptors, CD3, the IL-2

receptor, the growth hormone receptor, the insulin receptor, the acetylcholine receptor,

adrenergic receptors, VEGF receptors, and receptors for viruses such as the adenovirus receptor.

Aptamers useful according to the invention include but are not limited to aptamers

specific for the human MDR1 and MRP genes, including SEQ ID Nos: 67-117, SEQ ID Nos:

129-179 and 185-196, SEQ ID Nos: 199-235, SEQ ID Nos: 251-290, SEQ ID Nos: 293-384 of

WO 96/40715; aptamers specific for insulin receptor antibody, including SEQ ID Nos: 4-15 of

WO 95/30775; aptamers specific for peripheral blood mononuclear cells, including SEQ ID Nos:

7-39 of WO 96/34874; aptamers specific for multidrug resistance (MDR) phenotype, including

SEQ ID Nos: 88091 of WO 96/40715; aptamers specific for TNF $\alpha$ , including SEQ ID Nos: 209-

235 of WO 96/40717; and aptamers specific for cytokines, including those disclosed in WO

96/40717.

# Production of Nucleic Acid Molecules

- In one aspect, the invention pertains to nucleic acid molecules that can be used in the production of bifunctional nucleic acid molecules. In a preferred embodiment, these molecules are DNA or RNA molecules that can serve as templates for the assembly of bifunctional nucleic acid molecules, e.g. by possessing Watson-Crick complementarity to a bifunctional nucleic acid molecule and by directing the assembly of a complementary nucleic acid, e.g. by the action of a nucleotide polymerase. Such molecules are referred to herein as "bifunctional nucleic acid molecule-encoding sequences", and are useful for producing bifunctional nucleic acid molecules in which an aptamer and a biological effector sequence share a continuous backbone.
- Bifunctional nucleic acid molecule-encoding sequences can include sequences that are not Watson-Crick complementary to a bifunctional nucleic acid molecule, e.g. sequences that regulate and/or direct the activity of nucleotide polymerases, e.g. promoter sequences.
- Another aspect of the invention pertains to cloning vectors containing bifunctional nucleic acid molecules or bifunctional nucleic acid molecule-encoding sequences. As used herein, the term "cloning vector" refers to a nucleic acid molecule capable of transporting another nucleic acid molecule to which it has been linked. One type of vector is a "plasmid", which refers to a circular, double-stranded nucleic acid loop into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector, wherein additional nucleic acid segments can be ligated into the viral genome. Cloning vectors can include one or more nucleic acid sequences that can regulate biochemical processes by cis-acting mechanisms, e.g. promoters and/or binding sites for template-directed nucleotide polymerases. Examples of cloning vectors include pBR322 (Watson (1988) Gene 70:399-403), pUC19 (Yanisch-Perron et al (1985) Gene 33:103-19), and pBK-CMV and pBK-RSV (both available from Stratagene, La Jolla, CA).
- The recombinant cloning vectors of the invention comprise a nucleic acid of the invention in a form useful for producing larger quantities of a molecule of the invention than are available prior to operations on the vector in vivo or in vitro. For example, a bacterial cloning plasmid containing a bifunctional nucleic acid molecule sequence or a bifunctional nucleic acid molecule-encoding sequence can be used to amplify the number of copies of the sequence by introduction
- 240 production of bifunctional nucleic acid molecules. In a preferred embodiment, these molecules are DNA or RNA molecules that can serve as templates for the assembly of bifunctional nucleic acid molecules, e.g. by possessing Watson-Crick complementarity to a bifunctional nucleic acid molecule and by directing the assembly of a complementary nucleic acid, e.g. by the action of a nucleotide polymerase. Such molecules are referred to herein as "bifunctional nucleic acid molecule-encoding sequences", and are useful for producing bifunctional nucleic acid molecules in which an aptamer and a biological effector sequence share a continuous backbone.
- 245 Bifunctional nucleic acid molecule-encoding sequences can include sequences that are not Watson-Crick complementary to a bifunctional nucleic acid molecule, e.g. sequences that regulate and/or direct the activity of nucleotide polymerases, e.g. promoter sequences.
- Another aspect of the invention pertains to cloning vectors containing bifunctional nucleic acid molecules or bifunctional nucleic acid molecule-encoding sequences. As used herein, the term "cloning vector" refers to a nucleic acid molecule capable of transporting another nucleic acid molecule to which it has been linked. One type of vector is a "plasmid", which refers to a circular, double-stranded nucleic acid loop into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector, wherein additional nucleic acid segments can be ligated into the viral genome. Cloning vectors can include one or more nucleic acid sequences that can regulate biochemical processes by cis-acting mechanisms, e.g. promoters and/or binding sites for template-directed nucleotide polymerases. Examples of cloning vectors include pBR322 (Watson (1988) Gene 70:399-403), pUC19 (Yanisch-Perron et al (1985) Gene 33:103-19), and pBK-CMV and pBK-RSV (both available from Stratagene, La Jolla, CA).
- 250
- 255
- 260

- 265 of the plasmid into a suitable bacterial host. Another type of cloning vector according to the invention is a plasmid which comprises: 1) a template for a bifunctional nucleic acid molecule, which bifunctional nucleic acid molecule comprises an RNA aptamer and an RNA biological effector sequence linked via a phosphodiester bond; and 2) cis-acting sequences sufficient to allow in vitro transcription by a suitable RNA polymerase, e.g. T7 polymerase. Recombinant cloning vectors of the invention can also include vectors that lack a biological effector sequence and a biological effector-sequence encoding sequence but include: 1) an aptamer or an aptamer-encoding sequence and 2) a site, e.g. a multiple cloning site, which can facilitate the insertion of a biological effector sequence or a biological effector-sequence encoding sequence in such a manner that the resulting sequence is, or can serve as a template for, a bifunctional nucleic acid molecule in which the aptamer and the biological effector sequence are joined by a phosphodiester bond.
- 275 Numerous methods for the synthesis of nucleic acid sequences, including aptamers and biological effector sequences, are available and can be applied to the production of bifunctional nucleic acid molecules. For example, RNA molecules can be produced by in vitro transcription. In this technique, a gene encoding the desired RNA is cloned into a plasmid so that the gene is operably linked to a promoter for a DNA-dependent RNA polymerase, e.g., T7 RNA polymerase. The plasmid is then admixed, in a suitable buffer, with the RNA polymerase under conditions such that transcription occurs.
- 285 The polymerase chain reaction can be used to synthesize a specific DNA sequence (Mullis, U.S. Pat. 4,683,202). A double-stranded template sequence is admixed with a thermophilic DNA-dependent DNA polymerase, excess 5' and 3' primers, and excess free nucleotides in a suitable buffer. The mixture is subjected to repeated thermal cycling in a manner that allows melting and reannealing of complementary sequences and coordinated activation and deactivation of the polymerase.
- 290 Nucleic acid molecules can also be synthesized by machines that execute chemical polymerization of nucleotides. One such machine is produced by Applied Biosystems of Foster City, CA.
- Biological effector sequences useful according to the invention

- 295 Examples of nucleic acid sequences that can serve as biological effector sequences include sequences that encode useful polypeptides or useful polynucleotides, sequences that are "antisense" to nucleic acids that are present in the target cells, nucleic acid enzymes (e.g. ribozymes), nucleic acid sequences that can regulate biochemical processes by cis-acting mechanisms (e.g. -9-promoters or ribosome binding sites), and other useful nucleic acid sequences. A sequence is "antisense" if it is Watson-Crick complementary to and can inhibit the biological function of another nucleic acid molecule.
- 300 Examples of coding sequences that are useful according to the invention include, but are not limited to, sequences encoding the cystic fibrosis transmembrane regulator (Genbank Accession No. M28668), which is useful for treating cystic fibrosis; Factor VIII (Genbank Acc. No. E00527), which is useful for treating hemophilia; hemoglobin beta chain (Genbank Acc. No. V00497), which is useful for treating thalassemias; alpha-1-antitrypsin (AAT) (Genbank Acc. No. E00195), which is useful for treating AAT deficiency; insulin (Genbank Acc. No. J00265), which is useful for treating type 1 diabetes mellitus; TGF-beta (Genbank Acc. No. X02812 J05114), which is useful for treating inflammation and atherosclerosis; conditionally toxic polypeptides such as herpes simplex virus thymidine kinase (Genbank Acc. No. V00470), which is useful for treating malignant tumors; and antisense, ribozyme, and nucleic acid ligand molecules.
- 310 Examples of nucleic acid sequences that possess enzymatic activity and are useful according to the invention include, but are not limited to, ribozymes directed at mRNAs for proteins such as N-ras (Scherer et al (1997) J Biol Chem 272:14304-13) and c-fos (Scanlon et al (1994) Proc Natl Acad Sci USA 91:1123-27), which are useful for treating malignant tumors, and ribozymes directed at molecules involved in the life cycle of viruses such as HIV (Zhang et al (1996) Biochem Biophys Res Commun 229:466-71; Raillard and Joyce (1996) Biochemistry 35:11693-701; Michienza et al (1996) Proc Natl Acad Sci USA 93:7219-24), influenza A virus (Tang et al (1994) J Med Virol 42:385-95), and Tobacco Mosaic Virus (de Feyter et al (1996) Mol Gen Genet 250:329-38), which are useful for treating the respective viruses. Other examples of nucleic acid sequences that possess enzymatic activity and are useful according to the invention include, but are not limited to, hairpin ribozymes; hammerhead ribozymes; Group I introns; Group II introns; ribozymes directed at molecules involved in the life cycle of viruses
- 320

- 325 such as hepatitis B virus, hepatitis C virus, hepatitis D virus, hepatitis E virus, foot and mouth disease virus, mouse hepatitis virus, Moloney murine leukemia virus, bovine leukemia virus, influenza B virus, herpes viruses, lymphocytic choriomeningitis virus, potato leafroll virus, arenaviruses, potato virus YV, respiratory syncytial virus, retroviruses, rhabdoviruses, hemorrhagic fever agents, coronaviruses, rhinoviruses, myxoviruses, paramyxoviruses, and arboviruses; ribozymes directed at integrin mRNAs; RNase P; hepatitis D virus ribozymes; ribozymes directed at beta amyloid protein precursor mRNA; ribozymes directed at alpha actin; Tetrahymena ribozymes; ribozymes directed at phosphorothionate bonds; ribozymes directed at oncogenes such as H-ras, c-fos, c-myc, PML/RAR alpha bcr/abl, p53, AML1/MTG8, lck, fyn; ribozymes directed at thymidylate synthetase mRNA; ribozymes directed at mdr-1 mRNA; ribozymes directed at mdr-2 mRNA; ribozymes directed at components of telomerases; ribozymes directed at GAP-43 mRNA; ribozymes derived from chicory yellow mottle virus; ribozymes derived from arabis mosaic virus; ribozymes directed at amelogenin mRNA; ribozymes directed at serum amyloid A mRNA; ribozymes directed at osteopontin mRNA; polynucleotide kinase ribozymes; ribozymes directed at lipoxigenase mRNA; ribozymes directed at cytokine mRNAs; at ribozymes directed at NF-IL6 mRNAs; RNA ligase ribozymes; DNA ligase ribozymes; ribozymes directed at alkaline phosphatase mRNAs; Neurospora VS ribozymes; DNA enzymes such as DNA metalloenzyme DNA ligases; ribozymes directed at beta-glucuronidase mRNAs; ribozymes directed at plasminogen activator inhibitor mRNAs; self-alkylating ribozymes; ribozymes directed at glucose-regulated protein mRNAs; ribozymes directed at yeast ADE1 gene mRNA; ribozymes directed at insulin receptor substrate mRNAs; RNA cyclase ribozymes; ribozymes directed at glucokinase mRNAs; ribozymes directed at Nramp mRNAs; ribozymes directed at aberrant immunoglobulin light chain mRNAs; avocado sunblotch ribozymes; ribozymes directed at npt mRNAs; Pneumocystis carinii ribozymes; ribozymes directed at chloramphenicol acetyltransferase mRNAs; ribozymes directed at acetyl CoA carboxylase mRNAs; trans-splicing ribozymes; ribozymes directed at platelet-derived growth factor mRNAs; ribozymes directed at tumor necrosis factor alpha mRNAs; ribozymes directed at urokinase receptor mRNAs; ribozymes directed at atrial natriuretic factor mRNAs; ribozymes directed at pleiotrophin mRNAs; ribozymes directed at beta-2 microglobulin mRNAs; DNA-cleaving ribozymes; ribozymes directed at coliphages; Didymium ribozymes; chloroplast ribozymes; RNA
- 335
- 340
- 345
- 350

- 355 polymerase ribozymes; DNA polymerase ribozymes; ribozymes directed at *Drosophila* white ribozymes; ribozymes that hydrolyze phosphoric acid; ribozymes directed at calretinin mRNAs; ribozymes directed at VEGF mRNAs; integrase ribozymes; sun Y ribozymes; ribozymes directed at methyltransferase mRNAs; peptide cleaving ribozymes; peptide synthesizing ribozyme; ribozymes directed at prion mRNAs; ribozymes directed at inhibin mRNAs; self-copying ribozymes; ribozymes directed at alpha lactalbumin mRNAs; aminoacyl esterase mRNAs; tobacco ringspot virus ribozymes; barley yellow dwarf virus ribozymes; Chlamydomonas ribozymes; ribozymes that can specifically cleave single-stranded DNA; ribozymes directed at histone mRNAs, and ribozymes directed at glucan branching enzyme mRNAs.
- Examples of antisense nucleic acids that are useful according to the invention include, but are not limited to, molecules that are antisense to genes for K-ras (Alemany et al (1996) Cancer Gene Ther 3:296-301), c-kit (Yamanishi et al (1996) Jpn J Cancer Res 87:534-42), and c-myc (Leonetti et al (1996) J Natl Cancer Inst 88:419-29), useful for treating malignant tumors; HIV tat (Biasalo et al (1996) J Virol 70:2154-61), which is useful for treating which is useful for treating HIV infection; *Staphylococcus aureus* alpha-toxin (Kernodle et al (1997) Infect Immun 65:179-84), which is useful for treating *S. aureus* infection; TYL CV Rep protein (Bendahmane and Gronenborne (1997) Plant Mol Biol 33:351-57), which is useful for treating TYCLV infection; and Hepatitis C core protein (Moradpour et al (1996) Virology 222:51-63) which is useful for treating Hepatitis C infection. Other examples of antisense molecules useful according to the invention include those that have as their targets nucleic acids encoding gastrin, urokinase, Ha-ras, reverse transcriptase, HIV TAR, HIV gag, HIV rev, HIV nef, mycobacterial proteins, Tobacco Mosaic Virus proteins, Hepatitis B virus proteins, Hepatitis C virus proteins, Hepatitis D virus proteins, Hepatitis E virus proteins, Hepatitis G virus proteins, foot and mouth disease virus proteins, mouse hepatitis virus proteins, Moloney murine leukemia virus proteins, bovine leukemia virus proteins, influenza A virus proteins, influenza B virus proteins, herpes virus proteins, lymphocytic choriomeningitis virus proteins, potato leafroll virus proteins, arenavirus proteins, potato virus YN proteins, respiratory syncytial virus proteins, retrovirus proteins, rhadovirus proteins, hemorrhagic fever agent proteins, coronavirus proteins, rhinovirus proteins, myxovirus proteins, paramyxovirus proteins, and arbovirus proteins, integrins, RNase P, beta amyloid protein precursor, alpha actin, ribozymes directed at oncogenes such as c-fos, c-
- 360 tobacco ringspot virus ribozymes; barley yellow dwarf virus ribozymes; Chlamydomonas ribozymes; ribozymes directed at alpha lactalbumin mRNAs; aminoacyl esterase mRNAs; ribozymes directed at prion mRNAs; ribozymes directed at inhibin mRNAs; self-copying ribozymes; ribozymes directed at methyltransferase mRNAs; peptide cleaving ribozymes; peptide synthesizing ribozyme; ribozymes directed at VEGF mRNAs; integrase ribozymes; sun Y ribozymes; ribozymes directed at calretinin mRNAs; ribozymes that hydrolyze phosphoric acid; ribozymes directed at calretinin mRNAs; polymerase ribozymes; DNA polymerase ribozymes; ribozymes directed at *Drosophila* white
- 365 Gene Ther 3:296-301), c-kit (Yamanishi et al (1996) Jpn J Cancer Res 87:534-42), and c-myc (Leonetti et al (1996) J Natl Cancer Inst 88:419-29), useful for treating malignant tumors; HIV tat (Biasalo et al (1996) J Virol 70:2154-61), which is useful for treating which is useful for treating HIV infection; *Staphylococcus aureus* alpha-toxin (Kernodle et al (1997) Infect Immun 65:179-84), which is useful for treating *S. aureus* infection; TYL CV Rep protein (Bendahmane and Gronenborne (1997) Plant Mol Biol 33:351-57), which is useful for treating TYCLV infection; and Hepatitis C core protein (Moradpour et al (1996) Virology 222:51-63) which is useful for treating Hepatitis C infection. Other examples of antisense molecules useful according to the invention include those that have as their targets nucleic acids encoding gastrin, urokinase, Ha-ras, reverse transcriptase, HIV TAR, HIV gag, HIV rev, HIV nef, mycobacterial proteins, Tobacco Mosaic Virus proteins, Hepatitis B virus proteins, Hepatitis C virus proteins, Hepatitis D virus proteins, Hepatitis E virus proteins, Hepatitis G virus proteins, foot and mouth disease virus proteins, mouse hepatitis virus proteins, Moloney murine leukemia virus proteins, bovine leukemia virus proteins, influenza A virus proteins, influenza B virus proteins, herpes virus proteins, lymphocytic choriomeningitis virus proteins, potato leafroll virus proteins, arenavirus proteins, potato virus YN proteins, respiratory syncytial virus proteins, retrovirus proteins, rhadovirus proteins, hemorrhagic fever agent proteins, coronavirus proteins, rhinovirus proteins, myxovirus proteins, paramyxovirus proteins, and arbovirus proteins, integrins, RNase P, beta amyloid protein precursor, alpha actin, ribozymes directed at oncogenes such as c-fos, c-
- 370 and Gronenborne (1997) Plant Mol Biol 33:351-57), which is useful for treating TYCLV infection; and Hepatitis C core protein (Moradpour et al (1996) Virology 222:51-63) which is useful for treating Hepatitis C infection. Other examples of antisense molecules useful according to the invention include those that have as their targets nucleic acids encoding gastrin, urokinase, Ha-ras, reverse transcriptase, HIV TAR, HIV gag, HIV rev, HIV nef, mycobacterial proteins, Tobacco Mosaic Virus proteins, Hepatitis B virus proteins, Hepatitis C virus proteins, Hepatitis D virus proteins, Hepatitis E virus proteins, Hepatitis G virus proteins, foot and mouth disease virus proteins, mouse hepatitis virus proteins, Moloney murine leukemia virus proteins, bovine leukemia virus proteins, influenza A virus proteins, influenza B virus proteins, herpes virus proteins, lymphocytic choriomeningitis virus proteins, potato leafroll virus proteins, arenavirus proteins, potato virus YN proteins, respiratory syncytial virus proteins, retrovirus proteins, rhadovirus proteins, hemorrhagic fever agent proteins, coronavirus proteins, rhinovirus proteins, myxovirus proteins, paramyxovirus proteins, and arbovirus proteins, integrins, RNase P, beta amyloid protein precursor, alpha actin, ribozymes directed at oncogenes such as c-fos, c-
- 375 Tobacco Mosaic Virus proteins, Hepatitis B virus proteins, Hepatitis C virus proteins, Hepatitis D virus proteins, Hepatitis E virus proteins, Hepatitis G virus proteins, foot and mouth disease virus proteins, mouse hepatitis virus proteins, Moloney murine leukemia virus proteins, bovine leukemia virus proteins, influenza A virus proteins, influenza B virus proteins, herpes virus proteins, lymphocytic choriomeningitis virus proteins, potato leafroll virus proteins, arenavirus proteins, potato virus YN proteins, respiratory syncytial virus proteins, retrovirus proteins, rhadovirus proteins, hemorrhagic fever agent proteins, coronavirus proteins, rhinovirus proteins, myxovirus proteins, paramyxovirus proteins, and arbovirus proteins, integrins, RNase P, beta amyloid protein precursor, alpha actin, ribozymes directed at oncogenes such as c-fos, c-
- 380 proteins, myxovirus proteins, paramyxovirus proteins, and arbovirus proteins, integrins, RNase P, beta amyloid protein precursor, alpha actin, ribozymes directed at oncogenes such as c-fos, c-

- 385 myc, PML/RAR alpha bcr/abl, p53, AML1/MTG8, lck, fyn thymidylate synthetase, mdr-1, mdr-2, components of telomerase, GAP-43, chicory yellow mottle virus proteins, arabis mosaic virus proteins, amelogenin, serum amyloid A mRNAs, osteopontin, lipoxigenase, cytokines mRNAs, NF-IL6, DNA ligase ribozymes, alkaline phosphatase, beta-glucuronidase, plasminogen activator inhibitor, glucose-regulated proteins mRNAs, insulin receptor substrate mRNAs, glucokinase, Nramp, immunoglobulin light or heavy chains, T cell receptor chains, Pneumocystis carinii proteins, chloramphenicol acetyltransferase, acetyl CoA carboxylase, platelet-derived growth factor, tumor necrosis factor alpha, urokinase receptor, atrial natriuretic factor, pleiotrophin, beta-2 microglobulin, calcitonin, VEGF mRNAs, integrase, methyltransferase, prion proteins, inhibin, alpha lactalbumin, aminoacyl esterase, tobacco ringspot virus proteins, barley yellow dwarf virus proteins, Chlamydomonas proteins, histone proteins, glucan branching enzyme, bacterial proteins, fungal proteins, ticketissal proteins, proteins involved in autoimmune diseases, mRNAs, tRNAs, nucleoprotein RNAs, and snRNAs.
- 390
- 395

#### Assembly of Bifunctional Nucleic Acid Molecules

- 400 Aptamers and biological effector sequences can be linked using many methods. For example, if an aptamer and a biological effector sequence comprise mutually complementary sequences, they can be annealed to each other. The length of the mutually complementary sequences in each of the aptamer and biological effector sequence will be sufficiently long to permit the two molecules to form a hybrid that will not dissociate under the following conditions: at least 15°C, pH 7.4, 125mM NaCl in physiological compatible buffer. This length typically is in the range of 8-50 base pairs, preferably 10-20 base pairs, most preferably, 10-15 base pairs. In order to determine if the length of mutual complementarity is sufficient, a functional assay can be performed in which the hybrid molecule is used to transfect a target cell for which the aptamer-portion of the bifunctional nucleic acid is specific (in terms of its selectively recognizing a cell surface molecule on the target cell); if the biological effector sequence is detected in the cell (either its direct presence or detection of a product or activity encoded by the effector sequence) at a level that is at least 20% greater and preferably 50% or more than a comparative transfection using the biological effector sequence alone (i.e., without the Watson-Crick base-paired aptamer or with an aptamer that is not base-paired or
- 405
- 410



- covalently bonded to the effector sequence), then the length of mutual complementarity is considered sufficient according to the invention.
- Typically, the aptamer and biological effector fragments are admixed in a suitable buffer and heated to approximately 90°C. The mixture is then allowed to cool to a temperature that allows Watson-Crick base pairing between the complementary sequences. The temperature below which this occurs can be estimated by adding together the temperature values of each base of one of the complementary sequences, viz. 2°C for A or T and 4°C for G or C. If it is desired to produce a bifunctional nucleic acid molecule that comprises concatamers of aptamers and/or biological effector sequences formed by Watson-Crick base pairing, the ratio of each component in the bifunctional nucleic acid molecule can be manipulated by adjusting the molar ratios of the nucleic acid molecules in the annealing reaction. Furthermore, the average length of the concatamers can be manipulated by adding, in appropriate molar ratio, a "termination sequence". A termination sequence comprises a nucleotide sequence that allows incorporation into a concatamerized bifunctional nucleic acid molecule via Watson-Crick base pairing, but does not comprise a second sequence that would allow further concatamerization.
- Another method of assembling a bifunctional nucleic acid molecule involves admixing the aptamer and the biological effector sequence with a polycation, e.g. a polyamine, e.g. polylysine. This is particularly useful for assembling noncovalently linked DNA sequences, since the polyanionic charge of the DNA allows assembly via electrostatic attraction.
- Another approach to assembling bifunctional nucleic acid molecules exploits ligase enzymes to covalently link an aptamer and a biological effector sequence. Ligases catalyze the formation of a phosphodiester bond between two nucleic acid molecules. For example, T4 DNA ligase uses as its substrates duplex DNA or RNA, or DNA/RNA hybrids. T4 RNA ligase uses single stranded DNA or RNA. E. coli DNA ligase acts on duplex DNA molecules which have compatible cohesive ends.
- Assay for Gene Transfer
- The bifunctional nucleic acid is assayed for its ability to transfer genes into a target cell. For studies aimed at determining transfection efficiency, the biological effector sequence of the bifunctional nucleic acid contains a marker gene for firefly luciferase. For pharmaceutical

applications, the bifunctional nucleic acid contains a gene whose expression will have a beneficial therapeutic effect. The bifunctional nucleic acid is incubated with the cells. After incubation, the cells are lysed and are assayed for gene expression. In the case of the luciferase reporter, luciferin and ATP are added to lysed cells and the light emitted is measured with a luminometer.

445 Cells are harvested on the day of assay by centrifugation at 1200 rpm for 5 min at room temperature. The cell pellet is resuspended in phosphate buffered saline and re-centrifuged. This operation is performed twice. The cell pellet is then suspended in RPMI 1640 (Gibco Ltd.) to make up a suspension of  $2.7 \times 10^6$  cells per ml. The cells are then aliquoted into tubes and 0.75 ml of RPMI medium added, followed by 0.04-0.08 ml of 1 mM FP peptide and finally 0.25 ml of DNA solution. The transfection is then allowed to proceed by incubating the cells at 37°C for 4 h. After this time, the cells are harvested by centrifugation at 2000 rpm. The cells are then suspended in 1 ml of RPMI and re-centrifuged. Finally, the cells are suspended in 0.5 ml RPMI containing 0.1% foetal bovine serum. At this stage, if desired, the cells are electroporated at 300 V and 250  $\mu$ F using conventional electroporation.

455 Each 0.5 ml of transfected cell suspension is transferred to a well of a 12 well plastic culture plate containing 1.5 ml of RPMI 10% FBS. The original transfection tube is rinsed with a further 1 ml of medium and the wash transferred to the culture dish making a final volume of 3 ml. The culture plate is then incubated at 37°C for 24-72 h in an atmosphere of 5% CO<sub>2</sub>. The contents of each well in the culture dish are transferred to centrifuge tubes and the cells collected by centrifugation at 13,000 rpm. The pellet is resuspended in 0.12 ml of Lysis Buffer (100 mM sodium phosphate, pH 7.8; 8 mM MgCl<sub>2</sub>; 1 mM EDTA; 1% Triton X-100 and 15% glycerol) and agitated with a pipette. The lysate is centrifuged at 13,000 rpm for 1 minute and the supernatant collected. 80  $\mu$ l of the supernatant are transferred to a luminometer tube. The luciferase activity is then assayed using a Berthold Lumat L9501 luminometer. The assay buffer used is Lysis buffer containing 10 mM Luciferin and 100 mM ATP. Light produced by the luciferase is integrated over 4 sec and is described as relative light units (RLU). The data are converted to RLU/ml of lysate, RLU/cell or RLU/mg protein (protein concentration of the lysate having been determined in this case by the BioRAD Lowry assay).

470 Use of this type of pharmaceutical composition *in vivo* or *ex vivo* with nucleic acid containing a gene of physiological importance, such as replacement of a defective gene or an

additional potentially beneficial gene function, is expected to confer long term genetic modification of the cells and be effective in the treatment of disease.

For example, a patient that is subject to a viral or genetic disease may be treated in

accordance with the invention via *in vivo* or *ex vivo* methods. For example *in vivo* treatments, a bifunctional nucleic acid of the invention can be administered to the patient, preferably in a biologically compatible solution or a pharmaceutically acceptable vehicle, by ingestion, injection, inhalation or any number of other methods. The dosages administered will vary from patient to patient; a "therapeutically effective dose" will be determined by the level of enhancement of function of the transferred genetic material balanced against any risk or deleterious side effects. Monitoring levels of gene introduction, gene expression and/or the presence or levels of the encoded anti-viral protein will assist in selecting and adjusting the dosages administered. Generally, a composition including a bifunctional nucleic acid will be administered in a single dose in the range of 10 ng - 100 ug/kg body weight, preferably in the range of 100 ng - 10 ug/kg body weight, such that at least one copy of the therapeutic gene is delivered to each target cell. The therapeutic gene will, of course, be associated with appropriate regulatory sequences for expression of the gene in the target cell.

*Ex vivo* treatment is also contemplated within the present invention. Cell populations can be removed from the patient or otherwise provided, transduced with a therapeutic gene in accordance with the invention, then reintroduced into the patient. In general, *ex vivo* cell dosages will be determined according to the desired therapeutic effect balanced against any deleterious side-effects. Such dosages will usually be in the range of  $10^5$ - $10^8$  cells per patient, daily, weekly, or intermittently; preferably  $10^6$ - $10^7$  cells per patient.

A bifunctional nucleic acid according to the invention may be used to treat X-linked  $\gamma$ -globulinemia. The bifunctional nucleic acid will contain the Bruton's tyrosine kinase gene (Vetrie et al., 1993, Nature 361:226-233), which is carried on a 2.1 kb fragment delineated by the PvuI site at position (+33) and the HindIII site at position (+2126). The therapeutic gene may also encode a splice site and poly A tail, which may include portions of the human  $\beta$  globin locus splice and poly A signals; i.e., a BamHI XbaI 2.8 kb 3' splice/poly A flanking sequence containing exon 2 IVSII - exon 3 - - poly A sequences.

A bifunctional nucleic acid containing the Bruton's tyrosine kinase gene is assembled as described herein and used to treat X-linked  $\gamma$ -globulinemia by introducing the bifunctional nucleic

acid directly into a patient for *in vivo* gene therapy or contact the bifunctional nucleic acid with pre-B cells for *ex vivo* therapy, as described in Martensson et al.; Eur. Jour. Immunol. 1987, 17:1499; Okabe et al.; Eur. Jour. Immunol. 1992, 22:37; and Banerji et al.; Cell 33:729, 1983, and administering the transfected pre-B cells into a patient afflicted with X-linked  $\gamma$ -globulinemia. A bifunctional nucleic acid for treatment of X-linked  $\gamma$ -globulinemia will include a ligand for targeting of a preB cell. Such ligands are well-known in the art and will be specific for and capable of targeting one or more of the following cell surface markers: CD9, CD10, CD19, CD20, CD22, CD24, CD38, CD40, CD72, and CD74.

510 A bifunctional nucleic acid described herein also may be used for treatment of Gaucher's disease. Gaucher's disease stems from one of two different genetic mutations. Gaucher's type 1 is a CCG --> CAG mutation, which results in an Arg --> Gln substitution at position 119 of the  $\beta$ -glucocerebrosidase polypeptide (Graves, DNA 7:521, 1988). Gaucher's type 2 is a CTG -> CCG mutation, which results in a Leu --> Pro substitution at position 444 of the Z-glucocerebrosidase polypeptide (Tsui, NEJM 316:570, 1987). The presence of a  $\beta$ -glucocerebrosidase gene encoding a wild type polypeptide is believed to substantially correct Gaucher's disease. Therefore, a therapeutic bifunctional nucleic acid useful according to the invention includes the  $\beta$ -glucocerebrosidase gene, as described in Horowitz et al., 1989, Genomics 4:87-96, which is carried, as disclosed in Horowitz et al., on a 9722 base pair fragment extending from a BamHI site in exon 1 to an EcoRV site 31 to polyadenylation site. This fragment contains 11 exons and all intervening sequences, with translational start in exon 2. Sequences conferring position-independent and tissue-specific gene expression may be included in the construct and are carried on an 11.8 kb XhoI - SacI fragment from pIII.lyx construct as described in Bonifer et al., 1990, Euro. Mol. Biol. Org. Jour. 9:2843.

525 A bifunctional nucleic acid containing the  $\beta$ -glucocerebrosidase gene is assembled as described herein and used to treat Gaucher's disease by introducing the bifunctional nucleic acid directly into the host for *in vivo* treatment, or into isolated macrophages for *ex vivo* therapy, as described in Immunology and Cell Biology, 1993, Vol. 71, pages 75-78 and introducing the transfected macrophages into a patient afflicted with Gaucher's disease. Expression of the wild type transgene in a patient afflicted with Gaucher's disease should result in correction of the diseased state. The bifunctional nucleic acid will contain an aptamer that specifically targets a cell surface

530

antigen on a macrophage. Such aptamers may be easily prepared from procedures well-known in the art and described hereinabove, for example, aptamers having specificity for and capable of targeting one or more of the following cell surface markers: CD14, CD16, CD26, CD31, CDw32, CD36, CD45RO, CD45RB, CD63, CD71, CD74, CD23, CD25 and CD69.

The cells targeted for *in vivo* or *ex vivo* gene transfer in accordance with the invention include any cells to which the delivery of the therapeutic gene is desired. Such cells will bear a cell surface marker for which a corresponding specific ligand is available or can be prepared to allow for cell-specific targeting according to the invention. For example, cells of the immune system such as T-cells, B-cells, and macrophages, hematopoietic cells, and dendritic cells, each cell of which bears one or more well-known cell surface receptors having corresponding aptamers which may be selected for use as a targeting ligand in the bifunctional nucleic acid of the invention, depending upon the selected cell. Using established technologies, stem cells may be used for gene transfer after enrichment procedures (see, for example, European Patent Applications 0 455 482 and 0 451 611, which disclose methods for separating stem cells from a population of hematopoietic cells). Alternatively, unseparated hematopoietic cells and stem cell populations may be used as a target population for DNA transfer as described herein.

Compositions that Facilitate Use of Bifunctional Nucleic Acid Molecules

Yet another aspect of the invention features compositions including the molecules of the invention and another substance (or substances) that can facilitate the use of bifunctional nucleic acid molecules to influence a cell's biologic function. Such compositions can, for example, include substances that facilitate the uptake of exogenous nucleic acids by enhancing endocytosis, phagocytosis, potocytosis, micropinocytosis, and/or macropinocytosis (uptake enhancing agents). Uptake enhancing agents can include, for example, cytokines, e.g. M-CSF, PDGF, EGF, and HGF, as well as components of bacteria such as Shigella and Salmonella species. Other examples of uptake enhancing agents can include substances used in conventional transfection methods, such as transfection mediated by calcium phosphate.

Other examples of substances that can facilitate the use of a bifunctional nucleic acid molecule to introduce a biological effector sequence into a cell include substances that allow the internalized nucleic acid to escape from a subcellular vesicle ("escape agents"). Nucleic acid

565 molecules labeled for visualization, e.g. fluorescently labeled, appear in a punctate pattern if they are contained in subcellular vesicles. According to the invention, a substance is an escape agent if, when admixed with a cells and labeled nucleic acid molecules, the molecules can form a less punctate and more diffuse pattern than if the substance is omitted from the mixture. A substance can also be an escape agent according to the invention if, when admixed with cells and a nucleic acid template for a polypeptide or a polynucleotide, expression of the encoded molecule by the cell is at least 20% greater than if the substance is omitted.

570 Escape agents can include, e.g., adenovirus particles (Curiel et al. (1991) Proc. Natl. Acad. Sci. USA 88:8850; Christen et al. (1993) Proc. Natl. Acad. Sci. USA 90:2122-2126), portions of diphtheria toxin, including the transmembrane portion (Fisher and Wilson (1997) Biochem J 321:49-58), pertingolysin O (Gottschalk et al (1995) Gene Ther 2:498-503), peptides comprising sequences derived from influenza hemagglutinin (e.g., Plank et al (1994) J Biol Chem 269:12918-24), chloroquine (Guy et al (1995) Mol Biotechnol 3:237-48), and glycerol (Zauner et al (1997) Exp Cell Res 232:137-45). Another type of escape agent is an activator of protein kinase C, such as tetradecanoylphorbol 12, 13- acetate or 1, 2-dioctanoylglycerol (Reston et al (1991) Biochim Biophys Acta 1088:270-76). Other examples of escape agents include listeriolysin O, various phospholipases, and nuclease inhibitors.

580 Yet another example of a type of substance that can facilitate the use of bifunctional nucleic acid molecules to introduce a biological effector sequence into cells can include pharmaceutically/pharmacologically acceptable (i.e., applicable to mixing with pharmaceutical agents for in vitro testing such as for example for ex vivo uses which may require further testing or processing for use in mammals or humans), physiologically compatible (i.e., applicable for injection into an animal, particularly a human), and biologically compatible (i.e., acceptable for in vitro use on live cells but not necessarily acceptable, although not excluding acceptability, for in vivo use) carriers.

590 Bifunctional nucleic acids according to the invention may be delivered to cells using additional cell transfection enhancing agents, such as lipids (e.g., cationic lipids), peptides (e.g., cationic peptides) or synthetic polymers, for example, soluble DNA/polylysine complexes can be generated (Li et al., Biochem. J. 12, 1763 (1973)). Polylysine complexes tagged with asialoglycoprotein have been used to target DNA to hepatocytes in vitro (Wu and Wu, J. Biol. Chem.

262, 4429 (1987); U.S. Patent 5,166,320). Lactosylated polylysine (Midoux et al. (1993) Nuc. Acids Res. 21, 871-878) and galactosylated histones (Chen et al. (1994) Human Gene Therapy 5, 429-435) have been used to target plasmid DNA to cells bearing lectin receptors, and insulin conjugated to polylysine (Rosenkrantz et al. (1992) Exp. Cell Res. 199, 323-329) to cells bearing insulin receptors. Monoclonal antibodies have been used to target DNA to particular cell types (Machy et al. (1988) Proc. Natl. Acad. Sci. USA 85, 8027-8031; Trubetskoy et al. (1992) Bioconjugate Chem. 3, 323-27 and WO 91/17773 and WO 92/19287).

Peptides derived from the amino acid sequences of viral envelope proteins have been used in gene transfer when coadministered with polylysine DNA complexes (Plank et al. (1994) J. Biol. Chem. 269, 12918-24). Trubetskoy et al. (ibid) and Mack et al. ((1994) Am. J. Med. Sci. 307, 138-143) suggest that cocondensation of polylysine conjugates with cationic lipids can lead to improvement in gene transfer efficiency. WO 95/02698 used viral components to attempt to increase the efficiency of cationic lipid gene transfer.

615

**Use of the Invention**

In another aspect, the invention pertains to methods of introducing nucleic acid molecules into cells. Such methods can apply, for example, to prokaryotic or eukaryotic cells in vitro or in vivo. As used herein, "in vitro" refers to circumstances in which cells are harbored and/or cultivated under artificial conditions. "In vivo" refers to circumstances in which cells are located on or within an organism. In vitro methods of the invention can include admixture of compositions of the invention with cells into which the introduction of a biological effector sequence is desired. Such methods can include the use of substances that can facilitate the use of bifunctional nucleic acid molecules to introduce a biological effector sequence into cells, e.g. substances that can be included in compositions of the invention. Methods of the invention can include uses of such substances other than inclusion in a composition of the invention, for example application of the substance to cells prior to or following application of a composition of the invention.

610

The bifunctional nucleic acid molecule can be admixed with a suitable buffer, e.g. Tris-EDTA or phosphate-buffered saline. Typically, 0.1 µg-100 µg of bifunctional nucleic acid molecule in buffer is admixed with  $10^3$ - $10^7$  cells that bear the target of the bifunctional nucleic acid molecule's aptamer.

620 In some methods of the invention, bifunctional nucleic acid molecules are delivered to a cell via receptor-mediated transfer in combination with conventional transfection methods to achieve a synergistic effect. Examples of such methods follow:

1. *Transfection mediated by DEAE-dextran*: Naked nucleic acid can be introduced into cells

625 by forming a mixture of the nucleic acid and DEAE-dextran and incubating the mixture with the cells. A dimethylsulfoxide or chloroquine shock step can be added to increase the amount of nucleic acid uptake. DEAE-dextran transfection is only applicable to *in vitro* modification of cells and can be used to introduce nucleic acid transiently into cells but is not preferred for creating stably transfected cells. Thus, this method can be used for short term production of a gene product but is not a method of choice for long-term production of a gene product. Protocols for DEAE-dextran-mediated transfection can be found in *Current Protocols in Molecular Biology*, Ausubel, F.M. et al.

630 (e's.) Greene Publishing Associates, (1989), Section 9.2 and in *Molecular Cloning: A Laboratory Manual*, 2nd Edition. Sambrook et al. Cold Spring Harbor Laboratory Press, (1989), Sections 16.41-16.46 or other standard laboratory manuals.

2. *Electroporation*: Naked nucleic acid can also be introduced into cells by incubating the

635 cells and the nucleic acid together in an appropriate buffer and subjecting the cells to a high-voltage electric pulse. The efficiency with which nucleic acid is introduced into cells by electroporation is influenced by the strength of the applied field, the length of the electric pulse, the temperature, the conformation and concentration of the nucleic acid and the ionic

640 composition of the media. Electroporation can be used to stably (or transiently) transfect a wide variety of cell types and is only applicable to *in vitro* modification of cells. Protocols for electroporating cells can be found in *Current Protocols in Molecular Biology*, Ausubel, F.M. et al. (e's.) Greene Publishing Associates, (1989), Section 9.3 and in *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Sambrook et al. Cold Spring Harbor Laboratory Press, (1989), Sections 16.54-16.55 or other standard laboratory manuals.

3. *Liposome-mediated transfection ("lipofection")*: Naked nucleic acid can be introduced into

645 cells by mixing the nucleic acid with a liposome suspension containing cationic lipids. The nucleic acid/liposome complex is then incubated with cells. Liposome mediated transfection can be used to stably (or transiently) transfect cells in culture *in vitro*. Protocols can be found in *Current Protocols in Molecular Biology*, Ausubel, F.M. et al. (e's.) Greene Publishing Associates, (1989), Section 9.4



650 and other standard laboratory manuals. Additionally, gene delivery in vivo has been accomplished using liposomes. See for example Nicolau et al. (1987) *Enz. Meth.* 149:157-176; Wang and Huang (1987) *Proc. Natl. Acad. Sci. USA* 84:7851-7855; Brigham et al. (1989) *Am. J. Med. Sci.* 298:278; and Gould-Fogente et al. (1989) *Gene* 84:429-438.

655 4. *Transfection mediated by CaPO<sub>4</sub>*: Naked nucleic acid can be introduced into cells by forming a precipitate containing the nucleic acid and calcium phosphate. For example, a HEPES-buffered saline solution can be mixed with a solution containing calcium chloride and nucleic acid to form a precipitate and the precipitate is then incubated with cells. A glycerol or dimethyl sulfoxide shock step can be added to increase the amount of nucleic acid taken up by certain cells. CaPO<sub>4</sub>-mediated transfection can be used to stably (or transiently) transfect cells and is only applicable to *in vitro* modification of cells. Protocols for CaPO<sub>4</sub>-mediated transfection can be found in Current Protocols in Molecular Biology, Ausubel, F.M. et al. (eds.) Greene Publishing Associates, (1989), Section 9.1 and in Molecular Cloning: A Laboratory Manual, 2nd Edition. Sambrook et al. Cold Spring Harbor Laboratory Press, (1989), Sections 16.32-16.40 or other standard laboratory manuals.

660 Generally, when naked nucleic acid is introduced into cells in culture (e.g., by one of the transfection techniques described above) only a small fraction of cells (about 1 out of 10(5)) typically integrate the transfected nucleic acid into their genomes (i.e., the nucleic acid is maintained in the cell episomally). Thus, in order to identify cells which have taken up exogenous nucleic acid, it is advantageous to transfect nucleic acid encoding a selectable marker into the cell along with the nucleic acid(s) of interest. Preferred selectable markers include those which confer resistance to drugs such as G418, hygromycin and methotrexate. Selectable markers may be introduced on the same nucleic acid molecule as the gene(s) of interest or may be introduced on a separate molecule.

675 In vivo methods of the invention can include methods of administration of a composition of the invention to an organism that harbors cells into which introduction of a biological effector sequence is desired. The organism can be any plant or animal, but in a preferred embodiment is chosen from among the following: a human, a domestic animal, a cultivated plant. The targeted cells can be integral to the recipient organism or can, for example, be commensal microbes harbored by that organism. Routes of administration can include, e.g., intravenous, intraarterial, intrathecal, intraperitoneal, subcutaneous, intradermal, epicutaneous, oral, ocular, mucosal, or aerosol.

According to the invention, internalization of a nucleic acid by a cell can be detected in

680 several ways. For example, if the nucleic acid encodes a polypeptide or a polynucleotide, internalization can be indicated by an increase in synthesis of the encoded molecule by the cell. If the nucleic acid encodes or comprises an antisense sequence, a ribozyme, or a nucleic acid ligand, internalization can be indicated by a decrease in net synthesis of the target molecule by the cell or a decrease in activity of the target molecule.

685 In another method for detecting internalization of a nucleic acid by a cell, the nucleic acid is conjugated to a detectable marker, e.g. a radioactive marker, an enzyme such as alkaline phosphatase or horseradish peroxidase, or a fluorochrome such as fluorescein isothiocyanate or phycoerythrin. Methods for effecting such conjugation are well known to those skilled in the art. For example, the nucleic acid can be biotinylated and admixed with a streptavidin-linked marker. The conjugates are then admixed with the cells according to a method of the invention, extracellular nucleic acid removed, e.g. by denaturation or exonuclease treatment and washing, and the marker detected. If the nucleic acid is internalized, marker uptake will be at least 20% greater than when marker alone is used in the assay. The aptamer and biological effector of a bifunctional nucleic acid molecule can be detected separately in such an assay if each is conjugated to a different marker, e.g. fluorochromes with different spectral characteristics.

#### Dosage, Mode of Administration, and Pharmaceutical Formulations

In the methods of the invention, a composition of the invention is administered in vivo or applied in vitro in such a manner and amount and on such a schedule that a biological effector sequence is introduced into an appropriate number and type of target cells such that the desired effect is achieved. The specific conditions of a method can depend, for example, on the aptamer and the biological effector sequence, the nature of the target cells, and the effect desired.

700 Bifunctional nucleic acid molecules described herein may be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in or suspension in, liquid prior to infection can also be prepared. The preparation can also be emulsified, or the bifunctional nucleic acid molecules encapsulated in liposomes. The active ingredients are often mixed with carriers which are pharmaceutically acceptable and compatible with the active ingredient. The term "pharmaceutically acceptable carrier" refers to a carrier that does not cause an allergic reaction or other untoward effect in subjects to whom it is administered. A "biologically compatible carrier"

- 710 is one which is compatible with in vitro cellular transfection. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the formulation can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, and/or pH buffering agents.
- 715 Bifunctional nucleic acid molecules of the invention can be administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories, and in some cases, oral formulations or formulations suitable for distribution as aerosols. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutically grades of mannitol, lactose, starch magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25-70%.
- 725 The bifunctional nucleic acid molecules of the invention can be formulated into the compositions as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or with organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamine, histidine, procaine, and the like.
- 730 The bifunctional nucleic acid molecules are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or therapeutically effective. The quantity to be administered depends on the subject to be treated, including, e.g., the subject's body mass. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and may be peculiar to each subject. It will be apparent to those of skill in the art that the therapeutically effective amount of bifunctional nucleic acid molecules of this invention will
- 735

depend, *inter alia*, upon the administration schedule, the unit dose of aptamer and biological effectors sequence administered, whether the bifunctional nucleic acid molecules are administered in combination with other agents, the health of the recipient, and the therapeutic activity of the particular bifunctional nucleic acid molecule. In general, though, the dosage required for therapeutic efficacy will range from about 0.1 ug to 20 mg nucleic acid/kg body mass/dose.

Cells transfected with the bifunctional nucleic acid molecules of the invention can subsequently be administered to a subject to achieve a therapeutic effect, e.g. to provide a polypeptide which is encoded by the biological effector sequence and in which the subject is deficient. Typically, the cells will be formulated as a suspension in a pharmaceutically acceptable carrier, e.g. a physiologically compatible buffer. While the number of cells required for a therapeutic effect will vary according to both subject and cell characteristics, it will generally be between about  $10^7$  and  $10^9$  cells/dose.

## EXEMPLIFICATION

### Example 1

In one example, a cloning vector comprising a bifunctional nucleic acid molecule-encoding sequence is prepared in the following manner: The following oligodeoxynucleotides are synthesized: 1) 5'AACGGCGCGCGGCTAGTCCACACAGAACCCGTT3', the sense strand encoding a vascular endothelial growth factor-binding RNA (Jellinek et al, Biochemistry 1994 33:10450-10456) and 2) the complementary strand 5'AACGGCTTCTGTGCTGCTGACTAGCCGCGCGCGCTTCGA3' with an additional 3' Hind III compatible sequence. These oligonucleotides are admixed and annealed to each other. A nucleotide sequence including the E. coli beta galactosidase gene is prepared by digesting the pUC19 plasmid (Genbank accession number X02514) with Nar I and Hind III, and isolating the resulting 212 base pair fragment by agarose gel electrophoresis and elution. The annealed oligonucleotide and the pUC19-derived fragment are ligated using T4 DNA ligase. The resulting molecule is ligated to pSP70 plasmid (Promega, Madison, WI) that has been digested with Cla I and Xho I, exploiting the complementarity of Nar I and Cla I overhangs. The free Xho I end of the plasmid is blunted with Klenow and the plasmid is circularized with T4 DNA ligase. An

RNA bifunctional nucleic acid molecule is obtained by subjecting the plasmid to standard in vitro transcription procedures using SP6 RNA polymerase.

#### Example 2

In another example, the following oligodeoxynucleotides are synthesized:

1) 5'CGCGAAGCGCGCGGCTAGTCCACACAGAACCGTT3', the sense strand encoding a vascular endothelial growth factor-binding RNA (Jellinek et al, Biochemistry 1994 33:10450-10456) with an additional 5' Hae II compatible site and 2) the complementary strand 3'ACGCGTCTGTGTGTGACTAGCGCGCGCGTTCGGA3'. These oligonucleotides are admixed and annealed to each other. A nucleotide sequence including the E. coli beta galactosidase gene is prepared by digesting the pUC19 plasmid (Genbank accession number X02514) with Hae I and Hind III, and isolating the resulting 212 base pair fragment by agarose gel electrophoresis and elution. The annealed oligonucleotide and the pUC19-derived fragment are ligated using T4 DNA ligase. The resulting molecule is ligated to pSP70 plasmid (Promega) that has been digested with Hind III and Bgl II. The free Bgl II end of the plasmid is blunted with Klenow and the plasmid is circularized with T4 DNA ligase. An RNA bifunctional nucleic acid molecule is obtained by subjecting the plasmid to standard in vitro transcription procedures using SP6 RNA polymerase.

In an example of a method of the invention, the bifunctional nucleic acid molecules of Example 1 or Example 2 is introduced into CMS5 mouse fibrosarcoma cells that have been engineered to express a recombinant fusion polypeptide consisting of the intracellular and transmembrane portions of a fibroblast growth factor receptor (Genbank Acc. No. M34185) fused in-frame to the human VEGF 165 protein (see Swiss-Prot P15692), with the VEGF portion oriented extracellularly. About 0.1-100 ug of the bifunctional nucleic acid molecule is admixed in a suitable buffer with about  $10^3$  -  $10^7$  of these cells in vitro. Subsequent expression of beta galactosidase is determined by X-gal staining.

#### Example 4

795 A library of RNA's, each having 32 randomized bases between two fixed sequences of 30-45 bases, is constructed. An aptamer for the transferrin receptor (Genbank Acc. No. M11507) is isolated by SELEX, using the receptor's extracellular domain as a target. An antisense oligodeoxynucleotide which binds to the template region of the RNA component of human telomerase (5'TAGGGTTAG3') (Feng et al (1995) Science 269:1236-41) is synthesized and treated with calf alkaline phosphatase. The aptamer and the antisense biological effector sequence are ligated using T4 RNA ligase.

800 The resulting bifunctional nucleic acid is useful for treating malignant tumors. The bifunctional nucleic acid binds to transferrin receptor-bearing cells and delivers to the cell antisense oligonucleotide that binds to the RNA component of human telomerase, thus inhibiting function of telomerase enzyme.



835  $\times 10^6$  EGFP-expressing Jurkat cells in 500  $\mu$ L of a buffer consisting of 1 mM  $\text{CaCl}_2$ , 1 mM  $\text{MgCl}_2$ , 125 mM NaCl, 5 mM KCl, and 20 mM HEPES, pH 7.4 and incubation proceeded for 30 minutes at room temperature. 250  $\mu$ L of each suspension was diluted into duplicate tubes containing 750  $\mu$ L of RPMI 1640 media with 10% FCS and 100  $\mu$ M chloroquine.

840 Immediately after suspension in media, 250  $\mu$ L was removed and cells were centrifuged and washed twice in PBS buffer, suspended in 500  $\mu$ L PBS, and stained with propidium iodide for baseline fluorescence and viability analysis.

The remaining 750  $\mu$ L of cells and nucleic acid molecules were then incubated for 42

hours at 37°C with 95%  $\text{O}_2$ , 5%  $\text{CO}_2$ . Then another 250  $\mu$ L sample was drawn from each sample. These cells were then prepared and analyzed for fluorescence as at time zero.

The results of this experiment indicate that the aptamer-antisense conjugated molecules

845 exerted a greater antisense effect than antisense molecules alone. FACSscan analyses of viable

(determined by propidium iodide exclusion) Jurkat cells showed that, at 42 h, mean cell

fluorescence of cells incubated in a 0.5  $\mu$ M concentration of antisense alone was 13.4% below

that of appropriate control cells incubated with neither aptamer nor antisense. The mean

fluorescence of cells incubated with 0.5  $\mu$ M of the aptamer-antisense molecule was 25.7% below

850 that of appropriate controls incubated with 0.5  $\mu$ M of aptamer alone, so that at this concentration

the aptamer nearly doubled the effect of the antisense. Cells incubated in a 1  $\mu$ M concentration of

the aptamer-antisense combination had a mean fluorescence 23.7% below that of appropriate

controls, while the mean fluorescence of cells incubated with antisense alone was only 18.3%

below that of appropriate controls. Mean fluorescence of the two types of controls were

855 comparable, differing much less than 1%.

### Example 7

The aptamer and first 18 bases 5' of the antisense described in Example 6 are adjoined by

a covalent phosphodiester bond. This bifunctional molecule is a 98 base single stranded

oligodeoxynucleotide which has a sequence of:

860 5'-CTACCTACGATCTGACTAGCCGGAGCATGAGCGTTACGAAGTGCT

AAACGTAACGTTGCTTACTCTATGTTAGTTCCTGGTACCACTCGTCCCG3'



## Example 8

A bifunctional nucleic acid is constructed using the aptamer of Example 6 and an antisense molecule consisting of the 14 3' bases of the antisense molecule of Example 6 appended 3' to the phosphorothioate antisense molecule GEM 91, which inhibits the expression of HIV gag and is described in Veal et al, 1998, Antiviral Research 38:63-73. Aptamer and antisense are annealed in PBS at room temperature. The bifunctional nucleic acid molecule can target the antisense to T cells and is useful for treating HIV infection.

865

## Example 9

The reporter gene pEGFP-n2 from Clontech Laboratories is linearized by digestion with the restriction enzyme ApaI in a non-coding region leaving a 5' sense overhang with the sequence ACCGT on one end and an identical overhang on the 5' antisense strand. The following oligodeoxynucleotides are synthesized: 1)

870

5' TGCAGGGGGGGGGAACACTACATGAGAG 3', which has the first 4 bases at 5' end complementary to the ApaI overhang, an internal 8 bases which are complementary to the oligomer #2, and the final 15 bases which are Watson-Crick complementary to the 5' end of a 79 base human L-Selectin aptamer, #3, 2) 5' CCCCCCCCCC3' which is complementary to 8 bases internal in oligomer #1, and 3)

875

5' CTACCTACGATCTGACTAGCCGCGACATGAGCGTTACAAAGGTGCT  
AAACGTAACGTTGCTTACTCTATGTTCC3' which is a high affinity aptamer for human L-selectin (SEQ ID#134, Parma et al, PCT Publication WO 96/40703) The first two

880

oligodeoxynucleotides are annealed together by room temperature incubation and then are ligated to the linearized pEGFP-n2 plasmid with T4 DNA ligase. The linearized pEGFP-n2 plasmid now has two linkers ligated to both of its ends which are complementary to a 15 base region of the 5' end of the human L-selectin aptamer. The pEGFP-n2 plasmid is incubated with the

885

aptamer at a 1:2 molar ratio in a buffer consisting of 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 125mM NaCl, 5mM KCl, and 20mM HEPES, pH 7.4 for 30 minutes. Wild type Jurkat cells at a concentration of  $5 \times 10^5$  which endogenously express human L-selectin are then added to the plasmid and aptamer mix, and incubation continues for another 30 minutes. The mix is then equally divided and diluted into duplicate tubes containing RPMI 1640 with 10% FCS and 100µM chloroquine.

890

After 72 hours cells which have been stably transfected with pEGFP are selected for Neomycin resistance by addition of 1mg/mL of G418 antibiotic. Only cells which have stably integrated the neomycin resistance gene included in the pEGFP-n2 plasmid will survive.

A control group consists of cells which receive only linkers ligated to plasmid in a 2:1 molar ratio. For each group of cells a range of final pEGFP-n2 concentrations of 0 through 500 pM up to 1µM is used. The aptamer-plasmid molecule will yield a transfection efficiency at least 20% higher than that of plasmid alone, as measured by number of selected clones.

#### Example 10

A variation of example 6 would entail using the same human L-selectin aptamer oligodeoxynucleotide (SEQ ID#134, Parma et al, PCT Publication WO 96/40703) with a

sequence of:

5' CTACCTACGATCTGACTAGCCCGGACATGAGCGTTACAAAGGTCT

AAACGTAACGTTGCTTACTCTATGTTCC3' in conjunction with three distinct mixed

backbone antisense molecules. Each antisense molecule will retain the 18 phosphorothioate bases which are complementary to the EGFP start codon region, and will have at least one end with 14

phosphodiester bases complementary to the aptamer molecule. Antisense 1 (A1) has the sequence: 5'TGGTACCCACTCGTTCGCCGGATGGATGCTAGAC3' which is identical to the antisense

used in example 6 and consists of an EGFP hybridization region in the first 18 bases of the 5'

end and an aptamer hybridization region at the 14 bases of the 3' end. A2 has the sequence

5'AGAGTACATCAAGGTGGTACCCACTCGTTCGCC3' in which the first 14 bases at the 5'

end are complementary to the first 14 bases at the 3' end of the aptamer molecule, and the last 18 bases at the 3' end comprise the EGFP hybridization region. The A3 molecule contains aptamer

hybridization regions at both the 5' and 3' region and maintains the EGFP antisense sequence in

the internal 18 phosphorothioate base pairs: 5'

AGAGTACATCAAGGTGGTACCCACTCGTTCGCCGGATGGATGCTAGAC3'.

By varying the molar ratio of A1 and A2 to A3, it is possible to regulate the length of a

concatamerized aptamer-antisense chain. For example, aptamer, A1, A2, and A3 can be admixed in a molar ratio 3:1:1:2, so that the nucleic acid molecules thus formed will on average comprise three aptamer units to four antisense units. These molecules can then be used in the method

920 otherwise described in Example 6.

### EQUIVALENTS

Those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

## CLAIMS

1. A nucleic acid molecule comprising:  
a first nucleic acid sequence comprising an aptamer covalently linked to a second nucleic acid sequence comprising a biological effector sequence.
2. A nucleic acid molecule comprising:  
a first nucleic acid sequence comprising an aptamer linked via Watson-Crick base pairing to a second nucleic acid sequence comprising a biological effector sequence.
3. The molecule of claim 1 or 2, further comprising a third nucleic acid sequence which is an aptamer that is covalently linked to said nucleic acid molecule.
4. The molecule of claim 1 or 2, further comprising a third nucleic acid sequence which is an aptamer that is linked via Watson-Crick base pairing to said nucleic acid molecule.
5. The molecule of claim 3 wherein said third nucleic acid sequence comprises an aptamer that is different from said first nucleic acid sequence comprising an aptamer.
6. The molecule of claim 4 wherein said third nucleic acid sequence comprises an aptamer that is different from said first nucleic acid sequence comprising an aptamer.
7. The molecule of claim 1 or 2, comprising DNA and RNA.
8. The molecule of claim 1 or 2, wherein said biological effector sequence encodes a polypeptide or polynucleotide.
9. The molecule of claim 1 or 2, wherein said biological effector sequence comprises a messenger RNA.

- 945 10. The molecule of claim 8, wherein the coding sequence of said biological effector sequence comprises double-stranded DNA, and wherein said biological effector sequence comprises a promoter.
11. The molecule of claim 1 or 2, wherein said biological effector sequence comprises an antisense sequence.
- 950 12. The molecule of claim 1 or 2, wherein said biological effector sequence comprises a nucleic acid enzyme.
13. A nucleic acid molecule comprising a template for the assembly of the molecule of claim 1.
14. A cloning vector comprising the molecule of claim 1.
- 955 15. A cloning vector comprising the molecule of claim 11.
16. A composition comprising the molecule of claim 1 or 2 and a biologically acceptable carrier.
17. A composition comprising an admixture of a molecule of claim 1 or 2 and a cell that bears a target molecule for said aptamer.
- 960 18. A cell transfected with a nucleic acid molecule, wherein the nucleic acid molecule is chosen from the group: a molecule of claim 1 or 2, a molecule of claim 13, a vector of claim 14, and a vector of claim 15.
19. A method of introducing a biological effector sequence into a cell comprising contacting the molecule of claim 1 or 2 with a host cell.

- 965 20. A method of introducing a biological effector sequence into a cell using the molecule of claim 1 or 2, comprising administering said molecule to an organism.
21. The method of claim 20, which comprises administering to an organism a composition of claim 16.
- 970 22. A method of introducing a biological effector sequence into an organism, comprising introducing a biological effector sequence into a cell by contacting the molecule of claim 1 or 2 with a host cell, and administering said cell to an organism.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/15130

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6): A61K 31/70; C07H 21/02, 21/04; C12N 5/10, 15/85 US CL: Please See Extra Sheet According to International Patent Classification (IPC) or to both national classification and IPC		<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S.: 424/93.1; 435/320.1, 325; 514/44; 536/23.1, 23.2, 23.5, 24.1, 24.31		Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS: aptamer, nucleic acid ligand, antisense		<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>			
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	Y	US 5,270,163 A (GOLD et al.) 14 December 1993, see entire document, esp. col. 7, lines 12-42; col. 23, lines 39-58; col. 26, lines 60-64; col. 29, lines 11-16; col. 29, line 67-col. 30, line 16; col. 32, lines 49-57; col. 33, line 64-col. 34, line 13; col. 40, lines 23-32; col. 41, line 9; col. 43, lines 8-11; col. 52	Y	WO 92/14843 A1 (GILEAD SCIENCES, INC.) 03 September 1992, see entire document, esp. pages 26-29, 31, 32, 36-44, 48, 84, 87	Y	US 5,399,346 A (ANDERSON et al.) 21 March 1995, see entire document, esp. col. 2-3	Y	US 5,442,049 A (ANDERSON et al.) 15 August 1995, col. 4, at 11	Summary of the Invention
Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex. <input type="checkbox"/>											
Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *C* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is taken alone being obvious to a person skilled in the art *Z* document member of the same patent family											
Date of the actual completion of the international search 17 OCTOBER 1998		Date of mailing of the international search report 02 NOV 1998									
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230 Telephone No. (703) 308-0196 Authorized Officer ST PHILIP ZITOMER, PHD [Signature]											

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/15130

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/93.1; 435/20.1, 325; 514/44; 536/23.1, 23.2, 23.5, 24.1, 24.31